



CHECK OUT OUR MONTHLY PROMOTIONS ON:

TLRs • Inflammation • Dendritic Cell - T Cell Modulators • Host Defense

BRIDGING INNATE & ADAPTIVE IMMUNITY

The Journal of Immunology

This information is current as of October 25, 2010

How Immune Peptidases Change Specificity: Cathepsin G Gained Tryptic Function but Lost Efficiency during Primate Evolution

Wilfred W. Raymond, Neil N. Trivedi, Anastasia Makarova, Manisha Ray, Charles S. Craik and George H. Caughey

J. Immunol. 2010;185:5360-5368; originally published online Oct 1, 2010;

doi:10.4049/jimmunol.1002292

<http://www.jimmunol.org/cgi/content/full/185/9/5360>

References

This article **cites 61 articles**, 29 of which can be accessed free at: <http://www.jimmunol.org/cgi/content/full/185/9/5360#BIBL>

Subscriptions

Information about subscribing to *The Journal of Immunology* is online at <http://www.jimmunol.org/subscriptions/>

Permissions

Submit copyright permission requests at <http://www.aai.org/ji/copyright.html>

Email Alerts

Receive free email alerts when new articles cite this article. Sign up at <http://www.jimmunol.org/subscriptions/etoc.shtml>

The Journal of Immunology is published twice each month by The American Association of Immunologists, Inc., 9650 Rockville Pike, Bethesda, MD 20814-3994. Copyright ©2010 by The American Association of Immunologists, Inc. All rights reserved. Print ISSN: 0022-1767 Online ISSN: 1550-6606.



How Immune Peptidases Change Specificity: Cathepsin G Gained Tryptic Function but Lost Efficiency during Primate Evolution

Wilfred W. Raymond,* Neil N. Trivedi,^{†,‡} Anastasia Makarova,[‡] Manisha Ray,[§] Charles S. Craik,[§] and George H. Caughey*^{†,‡}

Cathepsin G is a major secreted serine peptidase of neutrophils and mast cells. Studies in *Ctsg*-null mice suggest that cathepsin G supports antimicrobial defenses but can injure host tissues. The human enzyme has an unusual “Janus-faced” ability to cleave peptides at basic (tryptic) as well as aromatic (chymotryptic) sites. Tryptic activity has been attributed to acidic Glu²²⁶ in the primary specificity pocket and underlies proposed important functions, such as activation of prourokinase. However, most mammals, including mice, substitute Ala²²⁶ for Glu²²⁶, suggesting that human tryptic activity may be anomalous. To test this hypothesis, human cathepsin G was compared with mouse wild-type and humanized active site mutants, revealing that mouse primary specificity is markedly narrower than that of human cathepsin G, with much greater Tyr activity and selectivity and near absence of tryptic activity. It also differs from human in resisting tryptic peptidase inhibitors (e.g., aprotinin), while favoring angiotensin destruction at Tyr⁴ over activation at Phe⁸. Ala²²⁶Glu mutants of mouse cathepsin G acquire tryptic activity and human ability to activate prourokinase. Phylogenetic analysis reveals that the Ala²²⁶Glu missense mutation appearing in primates 31–43 million years ago represented an apparently unprecedented way to create tryptic activity in a serine peptidase. We propose that tryptic activity is not an attribute of ancestral mammalian cathepsin G, which was primarily chymotryptic, and that primate-selective broadening of specificity opposed the general trend of increased specialization by immune peptidases and allowed acquisition of new functions. *The Journal of Immunology*, 2010, 185: 5360–5368.

Cathepsin G is a nonclassical endopeptidase of granulated immune cells. It is highly expressed in mast cells and neutrophils and to lesser extents in monocytes and dendritic cells (1, 2). In human mast cells, cathepsin G is abundant in secretory granules that contain histamine, chymase and tryptases (3, 4). In neutrophils, it resides with elastase in azurophil granules, where it is released into phagolysosomes and helps to kill microbes (5, 6). Cathepsin G and elastase also are released extracellularly, where they can resist antipeptidases, neutralize toxins, and kill bacteria (7, 8). High concentrations of uninhibited peptidase are present where neutrophils accumulate in large numbers, as in *Pseudomonas*-infected airways in humans with cystic fibrosis (9–11).

Cathepsin G belongs to a family of immune serine peptidases, with its closest relatives in humans being granzyme B and mast cell chymase. Genes for these enzymes arose by duplication and divergence of an ancestral chymase-like gene and are clustered on chromosome 14q11.2 (12, 13). Cathepsin G- and granzyme B-like genes separated early in evolution of placental mammals (13). Although phylogenetically related, these peptidases differ radically

in substrate specificity. Human granzyme B has caspase-like affinity for substrate Asp residues (14), and chymase is chymotryptic, hydrolyzing after Phe, Tyr, Trp, or Leu (15). Cathepsin G is less selective, with unusual combined ability to hydrolyze chymotryptic and tryptic (especially Lys-containing) substrates (16–18). Tryptic activity is thought to be responsible for activating proteinase-activated receptors (19), C3 (17), and prourokinase plasminogen activator (20). Despite its tryptic activity, cathepsin G has an active site that differs from known tryptic serine peptidases, most of which, like trypsin and tryptase, have a highly conserved “specificity triad” configuration of Asp¹⁸⁹, Gly²¹⁶, and Gly²²⁶ (chymotrypsinogen numbering). Crystal-derived structures of human cathepsin G (21) suggest that accommodation of basic side chains in tryptic substrates and inhibitors is due to Glu²²⁶ at the base of the primary specificity pocket. The side chain carboxylate of Glu²²⁶, which is nearly unique at this site among known peptidases, serves the charge–charge coupling function of Asp¹⁸⁹ in classic tryptic peptidases. In a similar fashion, with flip-flopping of charges, Arg²²⁶ determines specificity for acidic side chains (Asp-ase activity) in granzyme B (14). Although human cathepsin G activity is exceptionally broad, it is also comparatively weak, with specificity constants (k_{cat}/K_m) toward its best substrates being much lower than those of chymase (16, 22). Some functions may be independent of proteolytic activity; indeed, cathepsin G-derived peptides have antibacterial properties (23). Nonetheless, peptidase-dependent properties include secretagogue, angiotensin II-generating, metallopeptidase-activating, and hepatocyte growth factor-inactivating functions (24–28).

Studies in *Ctsg*^{-/-} mice suggest that cathepsin G is important for surviving fungal and bacterial infections, as from i.v. *Aspergillus fumigatus* (29) and *Staphylococcus aureus* (5). Immune

*Cardiovascular Research Institute, [†]Department of Medicine, and [§]Department of Pharmaceutical Chemistry, University of California at San Francisco, San Francisco, CA 94143; and [‡]Veterans Health Research Institute, San Francisco, CA 94121

Received for publication July 12, 2010. Accepted for publication August 31, 2010.

This work was supported in part by National Institutes of Health Grant HL024136 (to W.W.R. and G.H.C.) and a Veterans Affairs Career Development award (to N.N.T.).

Address correspondence and reprint requests to Dr. George H. Caughey at the current address: Veterans Affairs Medical Center 111-D, 4150 Clement Street, San Francisco, CA 94121. E-mail address: george.caughey@ucsf.edu

Abbreviations used in this paper: Ang, angiotensin; hCG, human cathepsin G; mCG, mouse cathepsin G; n, norleucine; 4NA, 4-nitroanilide; N.I., no inhibition; prouPA, prourokinase plasminogen activator; suc, succinyl.

deficits are even more pronounced when cathepsin G deficiency is combined with deficiency of neutrophil elastase and may be due to ineffective microbial killing by neutrophils (5). Cathepsin G-deficient mouse neutrophils also respond defectively to activation by immune complexes (30). However, compared with wild-type mice, *Ctsg*^{-/-} mice have lower bacterial counts in a model of bacterial bronchitis produced by tracheal inoculation with *Pseudomonas aeruginosa*-embedded beads (31), suggesting that cathepsin G interferes with airway defenses (although without a demonstrated effect on inflammation or mortality). In a noninfectious model of renal ischemia-reperfusion injury, *Ctsg*^{-/-} mice are protected from death and have less inflammation and tissue damage (32). Thus, mouse models suggest that cathepsin G contributes to antimicrobial defenses and to survival postinfection but can harm the host and actually increase mortality. Recent data reveal that dual chymase–cathepsin G inhibitors reduce pathology associated with allergic and neutrophilic inflammation (33, 34). Insights concerning contributions of human cathepsin G to host defense derive mostly from *in vitro* data. These studies suggest that the enzyme can contribute to host defense by cleaving bacterial virulence factors (35) or detract from host defense by inactivating antimicrobial collectins (36) and neutrophil chemokine receptors (11). Humans with Papillon-Lefevre syndrome, which is characterized by hyperkeratosis and destructive periodontitis, are deficient in active cathepsin G and other immune peptidases (37, 38) due to defects in dipeptidyl peptidase I, which is the major activator of cathepsin G and related peptidases from proenzyme forms (39, 40).

The present study was prompted by structural comparisons of mouse and human cathepsin G suggesting that the mouse enzyme differs from the “dual-specificity” tryptic–chymotryptic human enzyme in key specificity-determining residues, instead resembling chymotryptic mast cell chymases. These predictions regarding functions of the heretofore uncharacterized mouse enzyme were tested by comparing substrate preferences of recombinant wild-type and humanized mutant forms of mouse cathepsin G with those of human cathepsin G and by tracking evolution of primary specificity-determining amino acids in mammals from inferred ancestral forms. The findings suggest that mouse and ancestral mammalian cathepsin G, unlike the human enzyme, are purely chymotryptic and that a small number of active site mutations in primate ancestors of human cathepsin G dramatically altered activity, inhibitor sensitivity, and substrate specificity, causing, most notably, acquisition of trypsin-like ability to hydrolyze substrates after lysine residues and new targeting functions.

Materials and Methods

Phylogenetic analysis

Full amino acid sequences of cathepsin G not already published or annotated were obtained by data mining, including Basic Local Alignment Search Tool searches of high-throughput genome sequence and whole-genome shotgun databases at the National Center for Biotechnology Information (Bethesda, MD) using human and mouse cathepsin G genes and cDNAs as query sequences. Marmoset sequence (*Callithrix jacchus*, Contig 30.531) was obtained from a search of genomic sequence archived at the Washington University Genome Sequencing Center (St. Louis, MO). Previously unreported amino acid sequences of cathepsin G were predicted from genomic DNA using existing cathepsin G gene structures as a template following standard rules for placement of intron–exon boundaries. Cathepsin G amino acid sequences were aligned using Geneious software (Biomatters, Auckland, New Zealand). A likely evolutionary path was constructed from mutations in specificity triad codons using established branching dates for primate ancestors and presuming that mutations accumulated in a stepwise fashion. Specificity triad codons in ancestral mammalian cathepsin G were predicted by identifying the most parsimo-

nous path from putative ancestral sequence to known extant mammalian cathepsin G sequences.

Production of wild-type mouse cathepsin G

Mouse cathepsin G catalytic domain cDNA was PCR-amplified from Integrated Molecular Analysis of Genomes and their Expression Consortium (<http://image.hudsonalpha.org>) clone BC12513 with the addition of an overhang sequence to permit ligation-independent cloning into pIEx-3 (EMD Chemicals, Gibbstown, NJ). The original expressed construct containing an N-terminal GST tag was insoluble. Therefore, the GST sequence was removed from the plasmid by restriction digestion, and the signal sequence was ligated in-frame with the downstream enterokinase cleavage site using a linker. The resulting expression plasmid, after confirmation of sequence, was transfected into adherent Sf9 cells with Insect GeneJuice transfection reagent (Novagen). Serum-free conditioned medium supernatant was collected 72 h after transfection. Secreted recombinant procathepsin G was purified from supernatant by NaCl gradient elution from a 75 × 750 mm Heparin-5 PW column (Tosoh Bioscience, King of Prussia, PA) equilibrated in 20 mM Tris-HCl (pH 7.4) containing 50 mM NaCl and 2 mM CaCl₂, with procathepsin G eluting at a concentration of ~450 mM NaCl. Peak fractions were desalted and concentrated to ~0.05 ml with 5k NMWL centrifugal filters (Biomax) and incubated for 4 h at 25°C with 0.16 U enterokinase (Novagen), which then was removed on a Mono S 5/5 HR cation exchange column eluted with a gradient of NaCl. Activated cathepsin G eluted at a concentration of ~900 mM NaCl.

Production of humanized mutants of mouse cathepsin G

Specificity site mutants were generated from the modified pIEx-3 wild-type mouse procathepsin G plasmid using QuikChange II site-directed mutagenesis kits (Agilent Technologies, Santa Clara, CA). Single-site mutant Ala²²⁶Glu was constructed using QuikChange-designed primers (<http://www.stratagene.com/sdmdesigner>). Mutant plasmid sequence was verified in both directions. A Ser¹⁸⁹Ala/Ala²²⁶Glu double mutant then was generated using primers targeting the Ser¹⁸⁹ codon on the Ala²²⁶Glu plasmid. Primers were as follows: Ala²²⁶Glu forward 5'-CAACAATGGT AACCTC-CAG AGGTATTCAC CAAAATCCAG AG-3', reverse 5'-AAGCAACAAT GGTAACCTC CAGCTGTATT CACCAAAATC CAGAGCTT-3'; Ser¹⁸⁹Ala forward 5'-CCGAGAGAAA GGAAGGCTGC CTTTCAGGGGT G-3', reverse 5'-CACCCCTGAA GCGAGCTTC CTTTCTCTCG G-3'. Both mutant proteins were expressed, purified, and activated as for the wild-type recombinant enzyme.

Determination of active peptidase concentration

To determine the level of active peptidase, recombinant, purified wild-type mouse cathepsin G was titrated with human α_1 -proteinase inhibitor (α_1 -antitrypsin; EMD Chemicals) in PBS containing 0.01% Triton X-100 and 0.05% DMSO. Residual activity was measured by addition of cathepsin G substrate succinyl (suc)-L-Val-Pro-Phe-4-nitroanilide (4NA) (1 mM) and monitoring change in absorbance at 410 nm on a Synergy 2 microplate spectrophotometer (BioTek, Winooski, VT). The active concentration of α_1 -proteinase inhibitor was determined by titrating against human leukocyte elastase (Elastin Products, Owensville, MO), the active site concentration of which was measured by sp. act. assay (41). Glu²²⁶ mutants of mouse cathepsin G could not be titrated against α_1 -proteinase inhibitor or other common inhibitors because of nonstoichiometric inhibition. Instead, active enzyme concentration was based on protein content of purified preparations of highest sp. act. Human cathepsin G (MP Biomedicals, Solon, OH) active concentration was determined by assay with 1 mM suc-L-Val-Pro-Phe-4NA from sp. act. derived from published kinetic values of active site-titrated enzyme (42).

Determination of P1 substrate preferences

To probe the influence of the P1 residue at the site of hydrolysis, human (0.8 μ M) and recombinant mouse cathepsin G (0.03 μ M) were used to screen a P1-diverse library of tetrapeptide substrates divided into 20 wells, each containing a fixed P1 residue (standard amino acids plus norleucine, excluding Cys), as described in connection with human mast cell chymase (15). Each P1-fixed well contains an equimolar mixture of 8000 peptides representing all combinations of residues at P2, P3, and P4, with each peptide containing the fluorogenic leaving group 7-amino-4-carbamoylmethylcoumarin. The total number of peptides in the P1-diverse library is 160,000, each at a concentration of 31 nM. Assays were performed at 37°C in 150 mM Tris (pH 8), 150 mM NaCl, 0.01% Tween 20, and 1% DMSO. Hydrolysis was initiated by addition of enzyme

and monitored fluorometrically with excitation at 380 nm and emission at 450 nm.

Hydrolysis of synthetic substrates, angiotensin, casein, and prourokinase plasminogen activator

Chymotryptic and tryptic activity of the enzymes first were compared using peptidic, colorimetric substrates. Chymotryptic activity was assessed using N-terminally blocked peptidyl 4NAs with Phe in the P1 position at the site of hydrolysis. Standard assays monitored an increase in absorbance at 410 nm of 1 mM substrate at 37°C in PBS containing 0.01% Triton X-100 and 0.05% DMSO spectrophotometrically in 96-well plates. Turnover number k_{cat} and Michaelis constant K_m were calculated using Prism 5 software (GraphPad, La Jolla, CA) from substrate hydrolysis rates determined over a range of substrate concentrations bracketing K_m . For selected substrates, kinetic constants also were determined in high ionic strength conditions (0.45 M Tris-HCl [pH 8], 1.8 M NaCl, and 10% DMSO). Tryptic activity was assessed using *N*-carbobenzoyloxy-L-Lys-thiobenzylester (0.1 mM) at 37°C in PBS containing 0.01% Triton X-100 and 0.05% DMSO. Generation of free thiol by substrate hydrolysis was detected by inclusion of 0.1 mM 5, 5'-dithio-bis(2-nitrobenzoate) in the reaction mixture and monitoring the change in absorbance at 412 nm. Angiotensin-cleaving activity of wild-type mouse and human cathepsin G was compared with that of recombinant human mast cell chymase (produced as described in Ref. 43) by incubating enzymes with 19 μM angiotensin I for 40 min at 37°C in PBS containing 0.01% Triton X-100. Hydrolyzed fragments were detected by reverse-phase HPLC as in this laboratory's published work (44, 45). General proteinase activity of human chymase and wild-type and mutant cathepsin G was assessed by incubation of purified enzymes with bovine casein in PBS for 1 h at 37°C, followed by SDS-PAGE and staining with Coomassie brilliant blue. Human prourokinase plasminogen activator (prouPA; Landing Biotech, Newton, MA) activation by cathepsin G was detected via change in absorbance at 410 nm for 2 h at 37°C in PBS containing 0.01% Triton X-100, 0.05% DMSO, 10 mg/ml heparin (bovine lung; Sigma-Aldrich, St. Louis, MO), and 1 mM benzoyl-L-Val-Gly-Arg-4NA (Sigma-Aldrich), which is cleaved by active urokinase plasminogen activator but not by the forms of cathepsin G in this study.

Determination of inhibitor susceptibility

Inhibitory constants (K_i) for wild-type (Lys¹⁵) aprotinin with mouse cathepsin G mutants and human cathepsin G were determined by assaying hydrolysis of 0.037–1 mM suc-L-Val-Pro-Phe-4NA in 0.45 M Tris-HCl (pH 8) containing 1.8 M NaCl and 10% DMSO in the presence of 12.5 and 25 μM aprotinin for mouse mutants and 1 and 2 mM Lys¹⁵-aprotinin for human cathepsin G. Observed K_m in the presence of each inhibitor was derived using Prism 5 software (GraphPad) by nonlinear regression fit of the velocity versus substrate concentration curve. K_i was calculated from observed K_m (K_m observed), and K_m was obtained without inhibitor according to the equation $K_i = I / (K_m \text{ observed} / K_m - 1)$, where I is the inhibitor concentration.

Results

Prediction of mammalian cathepsin G protein sequence

Twenty mammalian sequences containing the complete catalytic domain were identified by in silico translation of genomic sequence. No sequences identifiable as cathepsin G were found in DNA from nonmammalian genomes. Accession numbers (<http://www.ncbi.nlm.nih.gov/genbank/>) for cathepsin G sequences identified and compared in this work (Fig. 1) are as follows: human NP_001902; chimpanzee (*Pan troglodytes*) XP_522810; gorilla (*Gorilla gorilla gorilla*) CABD01205422; rhesus (*Macaca mulatta*) XP_001114339; galago (*Otolemur garnettii*) AAQR01659072; mouse (*Mus musculus*) NP_031826; Norway rat (*Rattus norvegicus*) NP_001099511; kangaroo rat (*Dipodomys ordii*) ABRO01262720; dolphin (*Tursiops truncatus*) ABRN01151994; cattle (*Bos taurus*) XM_581980 (1) and XM_587026 (2); pig (*Sus scrofa*) XP_001926799; little brown bat (*Myotis lucifugus*) AAPE01464601; cat (*Felis catus*) ACBE01202053; dog (*Canis lupus familiaris*) AAEX02019056; giant panda (*Ailuropoda melanoleuca*) ACTA01131456; tenrec (*Echinops telfairi*) AAIY01076445; rocky hyrax (*Procavia capensis*) ABRQ01191013; armadillo (*Dasypus novemcinctus*) AAG-V020373271; and elephant (*Loxodonta africana*) AAGU03084767.

	189	*	216	226
Human	RERKAAF	KGDSGG	PLLCNN	VHAGIVSYGKSSGVP
Chimpanzee	---	S---	---	---
Gorilla	---	S---	---	---
Rhesus	---	S-R---	---	RP-I---
Marmoset	---	T-R---	V-R-A---	SQTAT-A---
Galago	---	Q-S-R---	V-S---	NAR-T-A---
Mouse	---	S-R---	V-S-Q---	SNN-N-A-K
Rat	---	S---	V-S-Q---	S-N-A---
Kangaroo Rat	---	S-R---	V-Q---	QN-T-A-K
Dolphin	---	S-L---	---	NRF-T-A---
Cattle 1	---	Q-S-L---	V-S-Q-V---	DRM-T-A---
Cattle 2	---	G-ST-L---	V-S-Q---	DRM-T-A---
Pig	---	S-L---	V-GR-Q---	NRQ-T-A---
Little Brown Bat	---	S-L---	MV-Q---	SR-T-G-Y---
Cat	---	SS-S-L---	I-E---	DRR-I-A---
Dog	---	AG-S-L---	V-G-Q---	DRR-T-A---
Giant Panda	---	PQ-S-M---	V-G-Q---	SR-T-A---
Hedgehog Tenrec	---	E-SV-L---	V-Q---	LN-I-A-K
Rocky Hyrax	---	SV-L---	V-H---	NRV-S-A---
Armadillo	---	N-S-R---	V-Q---	NIL-I-A-K
Elephant	---	Q-SV-L-A---	V-Q---	P-T-A-I---

FIGURE 1. Mammalian cathepsin G specificity triad sequences. Aligned portions of the catalytic domains (amino acids 185–220 using standard chymotrypsinogen numbering) bracket specificity-determining “triad” residues 189, 216, and 226 (shaded). Ser¹⁹⁵ involved in hydrolysis is marked with an asterisk. Hyphens indicate identity with human sequence. Human cathepsin G differs from consensus at two of three (Ala¹⁸⁹ and Glu²²⁶) specificity triad residues (58). Mouse cathepsin G and most other mammalian proteases match consensus sequence at these residues (59).

Human cathepsin G specificity triad is doubly mutated compared with that of ancestral cathepsin G

As shown by the mammalian sequences in Fig. 1, two of the three primary specificity-determining [“specificity triad” (46)] residues differ in human cathepsin G relative to those of nonprimate cathepsin Gs. In comparison with mouse cathepsin G specifically, human cathepsin G replaces Ser with Ala at position 189 and Ala with Glu at position 226. The third triad residue (Gly²¹⁶) is invariant. The mouse triad (Ser¹⁸⁹/Gly²¹⁶/Ala²²⁶) is typical in that it matches cathepsin G consensus sequence at all three positions. It also matches the human chymase triad, even though chymase is less closely related in overall sequence than human cathepsin G and belongs to a distinct clade distinct from mammalian cathepsin G (45). As shown in Fig. 2, residue 189 and 226 codons reveal origins of differences between humans and extant mammals in these specificity-determining amino acids. The Ser¹⁸⁹Ala change is absent in our closest mammalian relative, the chimpanzee, and thus occurred within the past 7 million years, which is when humans and chimpanzees are thought last to have shared a common ancestor (47). The Ser¹⁸⁹Ala conversion resulted from a single change (T → G) in the first base of the Ser¹⁸⁹ codon, which is otherwise highly conserved. The T → A change at the same site in marmoset DNA, which yields a conservative Ser¹⁸⁹Thr change, appears to have arisen independently after separation of ape and new world monkey lineages. The more dramatic Ala²²⁶Glu change occurred earlier in primate evolution than the Ser¹⁸⁹Ala change, because although the former mutation is observed in humans, chimpanzee, and old world monkey (rhesus macaque), it is absent from more distantly related primates (marmoset and galago) and from nonprimates. Compared to marmoset and galago, humans have a single change (C → A) in the second base of the Ala²²⁶ codon causing missense conversion to Glu. Thus, Ala²²⁶Glu conversion arose by point mutation in the primate lineage 31–43 million years ago, between the time humans last shared a common ancestry with old and new world monkeys, respectively (47). Mouse and rat cathepsin G have the same residues in positions 189 and 226 as most mammals, excepting humans, chimpanzees, and marmosets, as noted. However, the base in the wobble position for Ala²²⁶ has undergone an A → T synonymous change (Fig. 2). Mice and rats form the only the group of mammals with such




FIGURE 2. Evolution of specificity-determining Ala¹⁸⁹ and Glu²²⁶ in human cathepsin G. Codon sequence is shown below the corresponding residue. Mammalian consensus and predicted ancestral specificity-determining Ser¹⁸⁹ (TCT) and Ala²²⁶ (GCA) sequence differ from human at both amino acids and at two nucleotides. Nucleotides and amino acids differing from predicted ancestral sequence are indicated in boldface and italics. Nucleotide changes in dog, cat, giant panda, cattle 2, rat, and mouse codons are synonymous. The likely evolutionary path (based on stepwise accumulation of missense mutations, as indicated) predicts that Ala¹⁸⁹ and Glu²²⁶ mutations in humans arose up to 7 million years ago and between 31 and 43 million years ago, respectively.

a change, which thus appears to have arisen independently. Similarly, domestic dogs and cats have a synonymous, independent change in the Ser¹⁸⁹ codon. From the data summarized in Fig. 2, we conclude that ancestral mammalian cathepsin G had the following specificity triad amino acid sequence and codons: Ser¹⁸⁹ (TCT)/Gly²¹⁶ (GGA)/Ala²²⁶ (GCA). Thus, the human specificity triad, compared with the ancestral sequence, is modified more extensively than in any other identified mammal, with changes in amino acid sequence arising from two nucleotide mutations, each of which was introduced at a different time point in primate evolution.

Mouse cathepsin G is more active than human cathepsin G toward chymotryptic substrates and lacks tryptic activity

As shown in Fig. 3, the preferences of mouse cathepsin for substrate residues in the P1 position at the site of hydrolysis are markedly restricted compared with those of the human enzyme. Human cathepsin G has broad chymotryptic activity, cleaving after Trp, Phe, and Tyr, with little preference for Tyr versus Phe. It also has substantial Leu-ase and Met-ase activity, with little or no ability to cleave after P1 acidic residues (granzyme B-like Asp-ase activity) and small aliphatic residues (neutrophil elastase-like activity). Its most unique feature, however, is tryptic activity, which is as strong as its Tyr-cleaving chymotryptic activity and is largely Lys-specific. By contrast, the mouse enzyme lacks the tryptic, Leu-ase and Met-ase activity of the human enzyme, and even its chymotryptic profile is narrower, showing preference for Tyr over Phe and little inclination to cleave after Trp. Furthermore, the maximal activity of mouse cathepsin G is much higher than that of human cathepsin G. After normalization for differences in enzyme concentration used in the combinatorial assay, the P1 Tyr substrate-hydrolyzing activity is 19-fold greater for mouse versus human cathepsin G, even though maximal activity for both enzymes was achieved with substrates containing Tyr in




FIGURE 3. Comparison of mouse and human cathepsin G P1 substrate specificity. Purified mouse and human enzymes were profiled using a P1-diverse combinatorial library of fluorogenic tetrapeptide substrates containing all combinations of standard amino acids (except Cys) plus norleucine (n) at positions P2–P4, with each library fixed at P1. Individual assays assess overall ability to hydrolyze tetrapeptides containing a particular P1 amino acid with an equimolar mixture of 8000 peptides varying in 20 residues each in positions P2, P3, and P4 on the N-terminal side of cleavage site P1. To facilitate comparison between human and mouse enzyme, output in fluorescence units is normalized to the percentage of maximum output for each enzyme, which was achieved for both enzymes by the P1 Tyr well. The human enzyme has tryptic, chymotryptic, Leu-ase, and Met-ase activity, without Asp-ase or elastase-like activity, whereas mouse specificity is much narrower and mainly chymotryptic. Error bars show SD ($n = 3$).

the P1 position. As revealed in Figs. 3 and 4, wild-type human and mouse cathepsin G both cleave a variety of peptidyl-4NA chymotryptic substrates with Phe at the site of cleavage. Suc-L-Val-Pro-Phe-4-NA and suc-L-Phe-Pro-Phe-4-NA are the best identified substrates for both enzymes, suggesting that subsite P3 is similar for both enzymes. As shown in Table I, K_m for suc-L-Val-Pro-Phe-4-NA is substantially and significantly lower for mouse than for human cathepsin G at each of the two buffer conditions




FIGURE 4. Comparison of chymotryptic activity of mouse versus human cathepsin G. The graph compares hydrolytic rates using peptidyl-4NA substrates with P1 Phe, normalized for the concentration of active peptidase (nM). White and black bars represent activity of human and mouse cathepsin G, respectively. All of the substrates were studied in PBS at 1 mM. Substrates are as follows: suc-L-Val-Pro-Phe-4-NA (VPF), suc-L-Ala-Ala-Pro-Phe-4-NA (AAPF), suc-L-Ala-Glu-Pro-Phe-4-NA (AEPF), suc-L-Phe-Pro-Phe-4-NA (FPF), acetyl-L-Arg-Glu-Thr-Phe-4-NA (RETF), and benzoyl-L-Phe-4-NA (F). Results represent mean \pm SD of two independent experiments.

Table I. Comparison of kinetics of chymotryptic (*suc-L-Val-Pro-Phe-4NA-hydrolyzing*) activity

Enzyme	Buffer	k_{cat} (s^{-1}) ^a	K_m (mM) ^a	k_{cat}/K_m
Mouse cathepsin G	1.8 M NaCl	41 ± 2	0.12 ± 0.02	330
	PBS	33 ± 10	0.20 ± 0.12	170
Ala ²²⁶ Glu Mouse cathepsin G	1.8 M NaCl	22 ± 2	3.10 ± 0.35	7.1
	PBS	12 ± 2	0.63 ± 0.05	19
Ser ¹⁸⁹ Ala/Ala ²²⁶ Glu Mouse cathepsin G	1.8 M NaCl	16 ± 2	1.88 ± 0.29	8.5
	PBS	9 ± 0	0.37 ± 0.05	24
Human cathepsin G	1.8 M NaCl	100 ± 7	1.73 ± 0.19	58
	PBS ^b	14 ± 0	0.69 ± 0.12	20
Human chymase	1.8 M NaCl	230 ± 35	0.32 ± 0.12	720
	PBS ^b	17 ± 0	0.29 ± 0.07	59

^aMean ± SD, derived from three independent measurements at each of four substrate concentrations.

^bAs previously determined in Ref. 48.

($p = 0.0001$ and $p < 0.01$ in 1.8 M NaCl and PBS, respectively), possibly because the mouse enzyme better accommodates binding of the P1 Phe side chain, as suggested by modeling of the primary specificity pocket (data not shown). Neither enzyme shows much activity toward the human chymase-optimized substrate acetyl-L-Arg-Glu-Thr-Phe-4NA (48), indicating that the mouse enzyme, although it has a chymase-like specificity triad lining its primary (P1) specificity pocket, does not resemble chymase in its extended substrate binding fold. Although mouse cathepsin G is more active than human cathepsin G toward most of the surveyed chymotryptic substrates, the mouse enzyme is remarkable for reduced or absent tryptic activity, as reflected by minimal to no ability to cleave tetrapeptide substrates with basic amino acids in the P1 position (Fig. 3) and by lack of Lys-thioesterase activity (Fig. 5).

Humanized mutants of mouse cathepsin G acquire tryptic activity

As shown in Fig. 5, Lys-thioesterase activity appears in the humanized mutants of mouse cathepsin G, suggesting particularly that the Ala²²⁶Glu mutation contributes to tryptic activity in the human enzyme. Key modifications of the active site in human cathepsin G proposed to result in tryptic activity are modeled in Fig. 6.

Human but not wild-type mouse cathepsin G activates prouPA

As shown in Fig. 5, mouse cathepsin G has no detectable prouPA-activating activity, which requires tryptic cleavage at Lys¹⁵⁸. However, wild-type human cathepsin G and the humanized mutants possess this activity, suggesting that the human enzyme's ability to influence fibrinolysis was acquired with the active site Ala²²⁶Glu mutation.

Mouse cathepsin G with humanizing mutations acquires human-like sensitivity to Lys¹⁵-aprotinin

As shown in Fig. 5, wild-type mouse cathepsin G completely resists inactivation by Lys¹⁵-aprotinin, which is a broad-spectrum inhibitor of trypsin-like serine peptidases. However, wild-type human cathepsin G and humanized mouse mutants are sensitive, consistent with their tryptic activity, including hydrolysis of the Lys-thioester.

Mouse cathepsin G with humanizing mutations loses sensitivity to α_1 -proteinase inhibitor

Wild-type mouse cathepsin G is highly sensitive to α_1 -proteinase inhibitor, which, therefore, was used to titrate active sites in the enzyme preparations. However, the Glu²²⁶ mutants—and the wild-type human cathepsin G [consistent with prior reports (49)]—are resistant (data not shown).

Mouse cathepsin G preferentially destroys but human cathepsin G activates angiotensin I

As revealed by chromatograms in Fig. 7A, wild-type human chymase and human and mouse cathepsin G exhibit distinct patterns of angiotensin I hydrolysis. Recombinant human chymase, as reported (43), cleaves angiotensin I rapidly and highly selectively at Phe⁸ to generate vasoactive angiotensin II. Human cathepsin G also hydrolyzes selectively at Phe⁸, exhibiting very low level hydrolysis at Tyr⁴, but is much weaker than human chymase overall in attacking angiotensin I. In contrast, mouse cathepsin G preferentially hydrolyzes at Tyr⁴, which is an inactivating event in that it prevents angiotensin I from being activated to angiotensin II by hydrolysis at Phe⁸. Under these conditions, Phe⁸/Tyr⁴ selectivities for mouse cathepsin G, human cathepsin G, and human chymase are 0.31, 12, and 460, respectively. Although mouse cathepsin G hydrolyzes angiotensin at Tyr⁴ 3.2 times as often as at Phe⁸, it is 8.5-fold more active than human cathepsin G in generating angiotensin II, because it is more active overall, as also observed with chymotryptic 4NA substrates (Fig. 4, Table I). The mouse enzyme's preference for cleaving angiotensin at Tyr⁴ is consistent with its observed preference for tetrapeptide substrates with P1 Tyr versus Phe (Fig. 3).

Mouse and human cathepsin G are general proteinases

As shown by SDS-PAGE in Fig. 7B, wild-type mouse and human cathepsin G, as well as humanized mouse mutants, exhibit general caseinolytic activity. This contrasts with human mast cell chymase, which cleaves casein more selectively, suggesting that the general proteinase activity is not specifically a function of its specificity triad mutations, even though they broaden specificity.

Discussion

This work reveals that cathepsin G substrate specificity underwent major changes during evolution of primates. These changes broadened specificity to include one of human cathepsin G's most unusual characteristics (namely, tryptic activity) while reducing catalytic efficiency toward chymotryptic substrates. The phylogenetic analysis reveals that these transformations were generated by successive missense mutations in codons for key residues in the primary specificity pocket accommodating the substrate side chain at the site of hydrolysis. Consequently, as reflected by the identity of specificity triad residues 189 and 226, the human cathepsin G active site deviates substantially from that of the last ancestral cathepsin G shared by humans and new world monkeys, and indeed, its triad differs from that of any known serine peptidase. Mouse cathepsin G, which supports immune function as suggested by phenotypes of mice lacking cathepsin G (5, 29, 50), differs from the human enzyme in two of three triad residues.




FIGURE 5. Comparison of tryptic (Lys-containing) inhibitor and substrates in wild-type human cathepsin G (hCG), wild-type mouse cathepsin G (mCG), and humanized mutants Ala²²⁶Glu (A²²⁶E) and Ala²²⁶Glu/Ser189Ala (A²²⁶E/S¹⁸⁹A) mCG. The *left panel* compares strength of inhibition, as reflected by the ratio of K_m obtained in the presence (K_m observed) and absence (K_m) of 0.27 μ M Lys¹⁵-aprotinin, using the substrate suc-L-Val-Pro-Phe-4NA. K_i units are given in millimolar. The *middle panel* compares rates of hydrolysis of synthetic tryptic substrate carbobenzyloxy-L-Lys-thiobenzylester (100 μ M) normalized for peptidase concentration. The *right panel* compares rates of hydrolysis of human prouPA, which is activated by cleavage at Lys¹⁵⁸, as assessed by hydrolysis of urokinase plasminogen activator substrate suc-L-Val-Gly-Arg-4NA. N.I., no inhibition.

Instead, it has the predicted ancestral configuration. Thus, substrate preferences of the mouse enzyme, including lack of tryptic activity, more likely represent those of the ancestral form.

The mutagenesis results suggest that acquisition of tryptic activity was mainly due to the Ala²²⁶Glu change. Leaving aside the question of whether this change was beneficial, the tryptic anomaly in human cathepsin G is intriguing biochemically because it is, to our knowledge, a novel solution to the problem of generating tryptic specificity. The crux of that problem is to accommodate positively charged tips of basic amino acid side chains in the generally hydrophobic interior of the globular catalytic domain.





FIGURE 6. Specificity pocket mutations leading to acquisition of tryptic activity. The diagram depicts amino acids of the specificity triad lining the primary specificity pocket of cathepsin G and related serine peptidases, with examples of side chains of substrate P1 residues at the site of hydrolysis. The Ala²²⁶Glu mutation converted a chymase-like chymotryptic enzyme into a tryptic enzyme capable of hydrolyzing peptides after Lys, as shown, in part due to introduction of the negatively charged Glu side chain, which attracts the positive charge of the P1 Lys ϵ amino group. Configurations of classic chymotryptic and tryptic peptidases plus bovine duodenase, which uses a different configuration to accommodate tryptic and chymotryptic substrates, are shown for comparison. “Tryptase” refers to mast cell β -like tryptases, rather than to human α tryptase, which has an active site blocked by a Gly²¹⁶Asp mutation (60, 61).

serine peptidases (including immune peptidases) specialized to cleave substrates *exclusively* at tryptic sites, the specificity triad configuration is universally Asp¹⁸⁹/Gly²¹⁶/Gly²²⁶ (Fig. 6). In classic tryptic peptidases, the anionic Asp¹⁸⁹ side chain carboxylate at the base of the primary specificity pocket forms a charge-charge complex with the substrate Lys or Arg side chain. Probably the closest parallel with the human cathepsin G configuration is that of bovine duodenase, which also has dual tryptic and chymotryptic specificity but with specificity triad Asn¹⁸⁹/Gly²¹⁶/Asp²²⁶, with Asp²²⁶ proposed to serve the function of Asp¹⁸⁹ in classic tryptic serine peptidases (51). Duodenase is related to the chymase-cathepsin G-granzyme B family but forms its own clade, is absent in humans and mice, and is prominent in ruminant mammals, which also contain chymase and cathepsin G. The specificity triad of both types of cattle (*Bos taurus*) cathepsin G is identical to that of mouse and predicted ancestral cathepsin G and thus is expected to lack tryptic activity. The ancestors of duodenase thus achieved dual specificity independently, but because duodenase and cathepsin G are thought to share a chymotryptic ancestor (13, 46), the starting point was apparently the same. It is tempting to consider that duodenase in cattle plays a role similar to cathepsin G in humans, but not enough is known about roles and sources of these enzymes to inform speculation about functional parity. It may be significant that duodenase and cathepsin G (along with chymases and granzyme B-related peptidases) belong to the subset of immune peptidases that form only three disulfide linkages within the catalytic domain (13), the fewest such linkages known in the extended trypsin/chymotrypsin family. It is proposed that absence of particular disulfide pair (Cys¹⁹¹/Cys²²⁰) in the vicinity of the active site (and otherwise highly conserved in serine peptidases) allows structural plasticity (46, 52) such that changes in residues lining the pocket are better tolerated than in peptidases with the disulfide pair. This may explain how primate cathepsin G and perhaps duodenase were able to acquire tryptic activity with as little as a single amino acid change, even though changing chymotrypsin into a tryptic enzyme requires exchange of many more residues, including loops (53).

Although the mouse Ser¹⁸⁹Ala/Ala²²⁶Glu double mutant specificity triad matches human, the Ala²²⁶Glu single mutant specificity triad matches rhesus macaque (an old world monkey) and the great apes gorilla and chimpanzee. Because tryptic activity of

FIGURE 7. Peptidase and general proteinase activity. *A*, HPLC chromatograms of angiotensin (Ang) I after incubation with human chymase, human cathepsin G, and wild-type mouse cathepsin G. Peaks (as detected by monitoring absorbance at 210 nm) correspond to dipeptide HL and active octapeptide product Ang II (both resulting from hydrolysis at Phe⁸), inactive tetrapeptide DRVY and hexapeptide IHPFHL (resulting from hydrolysis at Tyr⁴), and uncleaved Ang I, as indicated. *B*, Results of SDS-PAGE of casein after incubation with peptidases mouse cathepsin G (mCG), Ala²²⁶Glu and Ser¹⁸⁹Ala/Ala²²⁶Glu mutants, and human cathepsin G (hCG). Size in kilodalton and elution positions of marker proteins are indicated.



the single mutant lies between that of the mouse wild-type and the double mutant, as reflected by Lys¹⁵-aprotinin K_i and by rates of Lys-thioester hydrolysis (Fig. 5), tryptic activity of macaque and chimpanzee cathepsin G may be intermediate between that of mouse and human wild-type enzymes. However, prouPA activation, which occurs at Lys¹⁵⁸ (20) is less robust in the double than the single mutant. We speculate that the Ser¹⁸⁹Ala mutation allows fuller expression of tryptic activity in human cathepsin G for some substrates because the Ala side chain is shorter, therefore partially offsetting the loss of space in the primary specificity pocket caused by the Glu for Ala exchange. Ala is also more hydrophobic than Ser and may increase hydrophobic interactions with the hydrophobic neck (rather than the charged tip) of substrate Lys side chains, thereby improving binding efficiency and K_m . Possibly, the Ser¹⁸⁹Ala mutation conferred an evolutionary advantage due to more efficient tryptic activity toward some substrates or more effective control by inhibition. Because neither of the humanized mouse cathepsin G mutations generated an enzyme with tryptic activity matching that of the human enzyme, additional changes in the vicinity of the active site affecting Lys substrate binding or turnover likely occurred to increase tryptic activity, perhaps under selective pressure. Potentially, other residues differing between mouse and human cathepsin G make as great a contribution as the Ala/Glu²²⁶ difference to the acquisition of tryptic specificity, although results of modeling and the duodenase example suggest that this is unlikely.

Broadening of specificity in primate cathepsin G was associated with a loss in catalytic efficiency toward chymotryptic substrates, as revealed by the ~19-fold difference between mouse and human cathepsin G in hydrolysis rates of pooled P1 Tyr tetrapeptides in the combinatorial assay and as shown with various P1 Phe 4NA substrates in Fig. 4. This may be inevitable, because ability to accommodate a variety of substrates likely means that the substrate binding site—and particularly the primary specificity pocket hosting the P1 residue at the site of hydrolysis—is no longer optimized for a particular side chain, resulting in looser binding and higher K_m for a given substrate. The importance of interactions involving P4–P2 residues N-terminal to P1, especially for the human enzyme, is illustrated by the observed large differences

in rates of hydrolysis of tetra- and tripeptidyl substrates with invariant P1 Phe. For both enzymes, this is also revealed by very low rates of hydrolysis of benzoyl-L-Phe-4NA, for which there are few contacts outside of the primary specificity pocket. These considerations likely explain why mouse cathepsin G, despite having a primary specificity pocket more similar to human chymase than to human cathepsin G, lacks the selectivity of angiotensin II-generating (Phe⁸-hydrolyzing) activity that is prominent in both human enzymes, for which surface topography, charge, and hydrophilicity outside of the primary specificity pocket are important determinants of angiotensin hydrolysis (44, 54). The finding that mouse cathepsin G preferentially destroys angiotensin I may explain why mast cell protease 4 chymase appears to be the major source of serine peptidase-derived angiotensin II-generating activity in uninflamed mouse tissues (43, 55). However, our in vitro assays show that mouse cathepsin G generates approximately one molecule of active angiotensin II for every three molecules of angiotensin I that it inactivates. In settings where cathepsin G is released in high concentrations, as in pneumonia, local generation of angiotensin II by cathepsin G may be important.

The comparisons of caseinolytic activity suggest that despite differences in ability to cleave peptidic substrates all of the tested forms of cathepsin G have general proteinase activity and may be less selective than human chymase, which cleaves casein [and albumin (15)] at fewer sites. It should be noted that the recent acquisition of tryptic activity via the Ala²²⁶Glu mutation, with retention of broad specificity, in a sense reverses the proposed general course of evolution of granule-associated immune peptidases of the chymase-cathepsin G-granzyme B family. All of these peptidases, despite their current specificities, are suggested to have evolved originally from tryptic peptidases (with classic triad Asp¹⁸⁹/Gly²¹⁶/Gly²²⁶) followed by despecialization (46), resulting in broad but weak activity. Subsequent mutations led to acquisition of specialized activity and higher catalytic efficiency. More generally, these results reveal how small changes in structure in this family of immune peptidases can cause large changes in function, as also occurred in mouse, rat, and hamster α -chymase (56, 57), which switched from chymotryptic to elastolytic activity, and in guinea pig α -chymase, which switched from chymotryptic

to Leu-ase activity (45). Although cathepsin G appears to be microbicidal in mice (6), in the context of chronic airway infection in humans with cystic fibrosis, cathepsin G interferes with the killing of certain bacteria (11). Potentially, differences in the nature of immune deficits in mice and humans with dipeptidyl peptidase I deficiency/Papillon-Lefevre syndrome are in part due to differences in specificity and immune roles of cathepsin G in the two species. Whether the changes in substrate preferences caused by specificity triad mutations in cathepsin G fundamentally altered host defense functions in humans remains to be determined. Similarly, the importance to homeostasis of mutation-enabled capabilities, such as modulation of fibrinolysis by activating proPA, will require further investigation and may be modeled inadequately by mice.

Disclosures

The authors have no financial conflicts of interest.

References

- Campbell, E. J., E. K. Silverman, and M. A. Campbell. 1989. Elastase and cathepsin G of human monocytes. Quantification of cellular content, release in response to stimuli, and heterogeneity in elastase-mediated proteolytic activity. *J. Immunol.* 143: 2961–2968.
- Stoeckle, C., V. Sommandas, E. Adamopoulou, K. Belisle, S. Schiekofe, A. Melms, E. Weber, C. Driessen, B. O. Boehm, E. Tolosa, and T. Burster. 2009. Cathepsin G is differentially expressed in primary human antigen-presenting cells. *Cell. Immunol.* 255: 41–45.
- Schechter, N. M., A. M. Irani, J. L. Sprows, J. Abernethy, B. Wintroub, and L. B. Schwartz. 1990. Identification of a cathepsin G-like proteinase in the MC₁TC type of human mast cell. *J. Immunol.* 145: 2652–2661.
- Benyon, R. C., J. A. Enciso, and A. D. Befus. 1993. Analysis of human skin mast cell proteins by two-dimensional gel electrophoresis. Identification of tryptase as a sialylated glycoprotein. *J. Immunol.* 151: 2699–2706.
- Reeves, E. P., H. Lu, H. L. Jacobs, C. G. Messina, S. Bolsover, G. Gabella, E. O. Potma, A. Warley, J. Roes, and A. W. Segal. 2002. Killing activity of neutrophils is mediated through activation of proteases by K⁺ flux. *Nature* 416: 291–297.
- Pham, C. T. 2006. Neutrophil serine proteases: specific regulators of inflammation. *Nat. Rev. Immunol.* 6: 541–550.
- Owen, C. A., M. A. Campbell, P. L. Sannes, S. S. Boukedes, and E. J. Campbell. 1995. Cell surface-bound elastase and cathepsin G on human neutrophils: a novel, non-oxidative mechanism by which neutrophils focus and preserve catalytic activity of serine proteinases. *J. Cell Biol.* 131: 775–789.
- Brinkmann, V., U. Reichard, C. Goosmann, B. Fauler, Y. Uhlemann, D. S. Weiss, Y. Weinrauch, and A. Zychlinsky. 2004. Neutrophil extracellular traps kill bacteria. *Science* 303: 1532–1535.
- Goldstein, W., and G. Döring. 1986. Lysosomal enzymes from polymorphonuclear leukocytes and proteinase inhibitors in patients with cystic fibrosis. *Am. Rev. Respir. Dis.* 134: 49–56.
- Liu, H., S. C. Lazarus, G. H. Caughey, and J. V. Fahy. 1999. Neutrophil elastase and elastase-rich cystic fibrosis sputum degranulate human eosinophils in vitro. *Am. J. Physiol.* 276: L28–L34.
- Hartl, D., P. Latzin, P. Hordijk, V. Marcos, C. Rudolph, M. Woischnik, S. Krauss-Etschmann, B. Koller, D. Reinhardt, A. A. Roscher, et al. 2007. Cleavage of CXCR1 on neutrophils disables bacterial killing in cystic fibrosis lung disease. *Nat. Med.* 13: 1423–1430.
- Caughey, G. H., T. H. Schaumberg, E. H. Zerweck, J. H. Butterfield, R. D. Hanson, G. A. Silverman, and T. J. Ley. 1993. The human mast cell chymase gene (CMA1): mapping to the cathepsin G/granzyme gene cluster and lineage-restricted expression. *Genomics* 15: 614–620.
- Gallwitz, M., J. M. Reimer, and L. Hellman. 2006. Expansion of the mast cell chymase locus over the past 200 million years of mammalian evolution. *Immunogenetics* 58: 655–669.
- Waugh, S. M., J. L. Harris, R. Fletterick, and C. S. Craik. 2000. The structure of the pro-apoptotic protease granzyme B reveals the molecular determinants of its specificity. *Nat. Struct. Biol.* 7: 762–765.
- Raymond, W. W., S. W. Ruggles, C. S. Craik, and G. H. Caughey. 2003. Albumin is a substrate of human chymase. Prediction by combinatorial peptide screening and development of a selective inhibitor based on the albumin cleavage site. *J. Biol. Chem.* 278: 34517–34524.
- Tanaka, T., Y. Minematsu, C. F. Reilly, J. Travis, and J. C. Powers. 1985. Human leukocyte cathepsin G. Subsite mapping with 4-nitroanilides, chemical modification, and effect of possible cofactors. *Biochemistry* 24: 2040–2047.
- Maison, C. M., C. L. Villiers, and M. G. Colomb. 1991. Proteolysis of C3 on U937 cell plasma membranes. Purification of cathepsin G. *J. Immunol.* 147: 921–926.
- Polanowska, J., I. Krokoszynska, H. Czapinska, W. Watorek, M. Dadlez, and J. Otlewski. 1998. Specificity of human cathepsin G. *Biochim. Biophys. Acta* 1386: 189–198.
- Sambrano, G. R., W. Huang, T. Faruqi, S. Mahrus, C. Craik, and S. R. Coughlin. 2000. Cathepsin G activates protease-activated receptor-4 in human platelets. *J. Biol. Chem.* 275: 6819–6823.
- Drag, B., and L. C. Petersen. 1994. Activation of pro-urokinase by cathepsin G in the presence of glucosaminoglycans. *Fibrinolysis* 8: 192–199.
- Hof, P., I. Mayr, R. Huber, E. Korzus, J. Potempa, J. Travis, J. C. Powers, and W. Bode. 1996. The 1.8 Å crystal structure of human cathepsin G in complex with Suc-Vac-Pro-PheP-(OPh)₂: a Janus-faced proteinase with two opposite specificities. *EMBO J.* 15: 5481–5491.
- Powers, J. C., T. Tanaka, J. W. Harper, Y. Minematsu, L. Barker, D. Lincoln, K. V. Crumley, J. E. Fraki, N. M. Schechter, G. G. Lazarus, et al. 1985. Mammalian chymotrypsin-like enzymes. Comparative reactivities of rat mast cell proteases, human and dog skin chymases, and human cathepsin G with peptide 4-nitroanilide substrates and with peptide chloromethyl ketone and sulfonyl fluoride inhibitors. *Biochemistry* 24: 2048–2058.
- Shafer, W. M., J. Pohl, V. C. Onunka, N. Bangalore, and J. Travis. 1991. Human lysosomal cathepsin G and granzyme B share a functionally conserved broad spectrum antibacterial peptide. *J. Biol. Chem.* 266: 112–116.
- Reilly, C. F., D. A. Tewksbury, N. M. Schechter, and J. Travis. 1982. Rapid conversion of angiotensin I to angiotensin II by neutrophil and mast cell proteinases. *J. Biol. Chem.* 257: 8619–8622.
- Sommerhoff, C. P., J. A. Nadel, C. B. Basbaum, and G. H. Caughey. 1990. Neutrophil elastase and cathepsin G stimulate secretion from cultured bovine airway gland serous cells. *J. Clin. Invest.* 85: 682–689.
- Owen, C. A., and E. J. Campbell. 1998. Angiotensin II generation at the cell surface of activated neutrophils: novel cathepsin G-mediated catalytic activity that is resistant to inhibition. *J. Immunol.* 160: 1436–1443.
- Shamamian, P., J. D. Schwartz, B. J. Pocock, S. Monea, D. Whiting, S. G. Marcus, and P. Mignatti. 2001. Activation of progelatinase A (MMP-2) by neutrophil elastase, cathepsin G, and proteinase-3: a role for inflammatory cells in tumor invasion and angiogenesis. *J. Cell. Physiol.* 189: 197–206.
- Raymond, W. W., A. C. Cruz, and G. H. Caughey. 2006. Mast cell and neutrophil peptidases attack an inactivation segment in hepatocyte growth factor to generate NK4-like antagonists. *J. Biol. Chem.* 281: 1489–1494.
- Tkalcevic, J., M. Novelli, M. Phylactides, J. P. Iredale, A. W. Segal, and J. Roes. 2000. Impaired immunity and enhanced resistance to endotoxin in the absence of neutrophil elastase and cathepsin G. *Immunology* 12: 201–210.
- Raptis, S. Z., S. D. Shapiro, P. M. Simmons, A. M. Cheng, and C. T. Pham. 2005. Serine protease cathepsin G regulates adhesion-dependent neutrophil effector functions by modulating integrin clustering. *Immunology* 22: 679–691.
- Sedor, J., L. Hogue, K. Akers, S. Boslaugh, J. Schreiber, and T. Ferkol. 2007. Cathepsin-G interferes with clearance of *Pseudomonas aeruginosa* from mouse lungs. *Pediatr. Res.* 61: 26–31.
- Shimoda, N., N. Fukazawa, K. Nonomura, and R. L. Fairchild. 2007. Cathepsin g is required for sustained inflammation and tissue injury after reperfusion of ischemic kidneys. *Am. J. Pathol.* 170: 930–940.
- Tausch, L., A. Henkel, U. Siemoneit, D. Poockel, N. Kather, L. Franke, B. Hofmann, G. Schneider, C. Angioni, G. Geisslinger, et al. 2009. Identification of human cathepsin G as a functional target of boswellic acids from the anti-inflammatory remedy frankincense. *J. Immunol.* 183: 3433–3442.
- Maryanoff, B. E., L. de Garavilla, M. N. Greco, B. J. Haertlein, G. I. Wells, P. Andrade-Gordon, and W. M. Abraham. 2010. Dual inhibition of cathepsin G and chymase is effective in animal models of pulmonary inflammation. *Am. J. Respir. Crit. Care Med.* 181: 247–253.
- López-Boado, Y. S., M. Espinola, S. Bahr, and A. Belaouaj. 2004. Neutrophil serine proteinases cleave bacterial flagellin, abrogating its host response-inducing activity. *J. Immunol.* 172: 509–515.
- Hirche, T. O., E. C. Crouch, M. Espinola, T. J. Brokelman, R. P. Mecham, N. DeSilva, J. Cooley, E. Remold-O'Donnell, and A. Belaouaj. 2004. Neutrophil serine proteinases inactivate surfactant protein D by cleaving within a conserved subregion of the carbohydrate recognition domain. *J. Biol. Chem.* 279: 27688–27698.
- de Haar, S. F., D. C. Jansen, T. Schoenmaker, H. De Vree, V. Everts, and W. Beertsen. 2004. Loss-of-function mutations in cathepsin C in two families with Papillon-Lefèvre syndrome are associated with deficiency of serine proteinases in PMNs. *Hum. Mutat.* 23: 524.
- Pham, C. T., J. L. Ivanovich, S. Z. Raptis, B. Zehnauer, and T. J. Ley. 2004. Papillon-Lefèvre syndrome: correlating the molecular, cellular, and clinical consequences of cathepsin C/dipeptidyl peptidase I deficiency in humans. *J. Immunol.* 173: 7277–7281.
- Wolters, P. J., C. T. Pham, D. J. Muilenburg, T. J. Ley, and G. H. Caughey. 2001. Dipeptidyl peptidase I is essential for activation of mast cell chymases, but not trypsinases, in mice. *J. Biol. Chem.* 276: 18551–18556.
- Adkison, A. M., S. Z. Raptis, D. G. Kelley, and C. T. Pham. 2002. Dipeptidyl peptidase I activates neutrophil-derived serine proteases and regulates the development of acute experimental arthritis. *J. Clin. Invest.* 109: 363–371.
- Nakajima, K., J. C. Powers, B. M. Ashe, and M. Zimmerman. 1979. Mapping the extended substrate binding site of cathepsin G and human leukocyte elastase. Studies with peptide substrates related to the alpha 1-protease inhibitor reactive site. *J. Biol. Chem.* 254: 4027–4032.
- Réhault, S., M. Brillard-Bourdet, M. A. Juliano, L. Juliano, F. Gauthier, and T. Moreau. 1999. New, sensitive fluorogenic substrates for human cathepsin G based on the sequence of serpin-reactive site loops. *J. Biol. Chem.* 274: 13810–13817.
- Caughey, G. H., W. W. Raymond, and P. J. Wolters. 2000. Angiotensin II generation by mast cell alpha- and beta-chymases. *Biochim. Biophys. Acta* 1480: 245–257.

44. Muilenburg, D. J., W. W. Raymond, P. J. Wolters, and G. H. Caughey. 2002. Lys40 but not Arg143 influences selectivity of angiotensin conversion by human alpha-chymase. *Biochim. Biophys. Acta* 1596: 346–356.
45. Caughey, G. H., J. Beauchamp, D. Schlatter, W. W. Raymond, N. N. Trivedi, D. Banner, H. Mauser, and J. Fingerle. 2008. Guinea pig chymase is leucine-specific: a novel example of functional plasticity in the chymase/granzyme family of serine peptidases. *J. Biol. Chem.* 283: 13943–13951.
46. Wouters, M. A., K. Liu, P. Riek, and A. Husain. 2003. A despecialization step underlying evolution of a family of serine proteases. *Mol. Cell* 12: 343–354.
47. Steiper, M. E., and N. M. Young. 2006. Primate molecular divergence dates. *Mol. Phylogenet. Evol.* 41: 384–394.
48. Raymond, W. W., S. Su, A. Makarova, T. M. Wilson, M. C. Carter, D. D. Metcalfe, and G. H. Caughey. 2009. Alpha 2-macroglobulin capture allows detection of mast cell chymase in serum and creates a reservoir of angiotensin II-generating activity. *J. Immunol.* 182: 5770–5777.
49. Travis, J., J. Bowen, and R. Baugh. 1978. Human alpha-1-antichymotrypsin: interaction with chymotrypsin-like proteinases. *Biochemistry* 17: 5651–5656.
50. Raptis, S. Z., and C. T. Pham. 2005. Neutrophil-derived serine proteases in immune complex-mediated diseases. *Immunol. Res.* 32: 211–215.
51. Pletnev, V. Z., T. S. Zanolodchikova, W. A. Pangborn, and W. L. Duax. 2000. Crystal structure of bovine duodenase, a serine protease, with dual trypsin and chymotrypsin-like specificities. *Proteins* 41: 8–16.
52. Zanolodchikova, T. S., E. V. Smirnova, A. N. Andrianov, I. V. Kashparov, O. D. Kotsareva, E. A. Sokolova, K. B. Ignatov, and A. D. Pemberton. 2005. Cloning and molecular modeling of duodenase with respect to evolution of substrate specificity within mammalian serine proteases that have lost a conserved active-site disulfide bond. *Biochemistry (Mosc.)* 70: 672–684.
53. Hedstrom, L., L. Szilagyi, and W. J. Rutter. 1992. Converting trypsin to chymotrypsin: the role of surface loops. *Science* 255: 1249–1253.
54. Sanker, S., U. M. Chandrasekharan, D. Wilk, M. J. Glynnias, S. S. Karnik, and A. Husain. 1997. Distinct multisite synergistic interactions determine substrate specificities of human chymase and rat chymase-1 for angiotensin II formation and degradation. *J. Biol. Chem.* 272: 2963–2968.
55. Lundequist, A., E. Tchougounova, M. Abrink, and G. Pejler. 2004. Cooperation between mast cell carboxypeptidase A and the chymase mouse mast cell protease 4 in the formation and degradation of angiotensin II. *J. Biol. Chem.* 279: 32339–32344.
56. Kunori, Y., M. Koizumi, T. Masegi, H. Kasai, H. Kawabata, Y. Yamazaki, and A. Fukamizu. 2002. Rodent alpha-chymases are elastase-like proteases. *Eur. J. Biochem.* 269: 5921–5930.
57. Kervinen, J., M. Abad, C. Crysler, M. Kolpak, A. D. Mahan, J. A. Masucci, S. Bayoumy, M. D. Cummings, X. Yao, M. Olson, et al. 2008. Structural basis for elastolytic substrate specificity in rodent alpha-chymases. *J. Biol. Chem.* 283: 427–436.
58. Salvesen, G., D. Farley, J. Shuman, A. Przybyla, C. Reilly, and J. Travis. 1987. Molecular cloning of human cathepsin G: structural similarity to mast cell and cytotoxic T lymphocyte proteinases. *Biochemistry* 26: 2289–2293.
59. Heusel, J. W., E. M. Scarpati, N. A. Jenkins, D. J. Gilbert, N. G. Copeland, S. D. Shapiro, and T. J. Ley. 1993. Molecular cloning, chromosomal location, and tissue-specific expression of the murine cathepsin G gene. *Blood* 81: 1614–1623.
60. Huang, C., L. Li, S. A. Krilis, K. Chanasyk, Y. Tang, Z. Li, J. E. Hunt, and R. L. Stevens. 1999. Human tryptases alpha and beta/II are functionally distinct due, in part, to a single amino acid difference in one of the surface loops that forms the substrate-binding cleft. *J. Biol. Chem.* 274: 19670–19676.
61. Marquardt, U., F. Zettl, R. Huber, W. Bode, and C. Sommerhoff. 2002. The crystal structure of human α_1 -tryptase reveals a blocked substrate-binding region. *J. Mol. Biol.* 321: 491–502.